

PREPARATION SHEET - BED BUGS

It is necessary for you to remove all pets, cover fish tanks and remain out of the unit for approximately 4 – 6 hours following the treatment. If you are pregnant or have any respiratory or medical issues it is recommended that you stay out of the unit for 24 hours.

A complete bedbug treatment is considered 2 full treatments 2 weeks apart. It is imperative that the 2nd treatment be completed in order for the treatment to be effective. When treating for bedbugs it is necessary to treat both the **bedroom** and **living room** areas.


BEDROOM PREPARATION




Remove and bag all items off floors and away from the walls. Vacuum all floor areas including around all baseboards and all furniture. All items must be removed from the dressers and nightstands.



All clothing items must be placed in a sealed bag. All other items should be placed in a sealed bag. Remove all items from the base of the closets and place in sealed bag.



All mattresses and box springs must be placed upright against the wall. All furniture should be moved at least 12 inches away from the walls.




Place all treated sealed bags in an area that do not require treatment.


****All items should remain in the dryer for approximately 45 minutes.****

LIVING ROOM PREPARATION


GENERAL PREPARATION




Remove all clutter off floors and away from the walls. Vacuum all floor areas including around all baseboards and all furniture.




All electrical wall plates should be removed.



All furniture should be moved at least 12 inches away from the walls.



All paintings, mirrors, and wall accessories should be removed from living room and bedroom walls. Gather all drapes, blinds, etc, and either tie back or fully open.



Place sofa on its side so technician can have access to the under area of the sofa.

AFTER TREATMENT INFORMATION:

Following the treatment, do not vacuum or wash within 6 inches of the baseboard area for approximately 3 weeks. Expect to continue to have activity for approximately **2 - 3 weeks** following the treatment.